

Optimal Signal Transduction

Gabriele Scheler

Carl Correns Foundation for Mathematical Biology
1030 Judson Drive
Mountain View, Ca. 94040

October 24, 2012

Abstract. We propose a model of parameter learning for signal transduction, where the objective function is defined by signal transmission efficiency. This is a novel approach compared to the usual technique of adjusting parameters only on the basis of experimental data. We apply this to learn kinetic rates as a form of evolutionary learning, and find parameters which satisfy the objective. These may be intersected with parameter sets generated from experimental time-series data to further constrain a signal transduction model. The resulting model is self-regulating, i.e. perturbations in protein concentrations or changes in extracellular signaling will automatically lead to adaptation. We systematically perturb protein concentrations and observe the response of the system. We find fits with common observations of compensatory or co-regulation of protein expression levels in cellular systems (e.g. PDE3, AC5). In a novel experiment, we alter the distribution of extracellular signaling, and observe adaptation based on optimizing signal transmission. Self-regulating systems may be predictive of unwanted drug interference effects, since they aim to mimic complex cellular adaptation in a unified way.

1 A transfer function approach to signal transduction

Signal transduction systems are usually modeled as networks of biochemical kinetic equations implemented as continuous-time dynamical models using differential equations [1]. If we regard a subset of species as inputs, and make sure that the system always converges to equilibrium values by using weakly reversible equations, we may transform these models into a set of matrices fulfilling the role of input-output transfer functions, i.e. a mapping from rectangular input signals of sufficient duration to steady-state concentrations for all target species [7]. We may then ask the question of optimal or efficient signal transmission for a variety of input scenarios, focusing on selected target species as relevant outputs or actuators of the system. An experimental analysis has shown "fold-change" responses to changes in input [5]. A theoretical analysis has shown that input-output transfer functions follow the shape of hyperbolic (saturating) curves, which are equivalent to sigmoids for logarithmically scaled input [7]. We hypothesize that an efficient signal transmission would maximize the response coefficient $R_{C,S}$ (the response of species C to input S) defined as

$$R_{C,S} = \frac{\frac{C_t}{C_0} - 1}{\log(\frac{S_t}{S_0}) - 1}$$

with concentration change of target C and input S from baseline ($t=0$) to signal time t . (We have used this both with and without $\log$, and report on logarithmically scaled input here, since the results seemed to be more biologically relevant.) We may also optimize for the slope s of the sigmoid at half-maximum concentration. This is equivalent to maximizing $R_{C,S}$, provided that the input signal remains entirely between the upper and lower boundaries of the sigmoid. However, shifting the sigmoid function to the left or to the right is even more important than slope in experimental settings. E.g., Fig. 1 A,B show the effect of a KO for the protein RGS in an experimental assay in yeast [9], and in our model. In addition to optimizing for signal transmission efficiency, we discuss




Figure 1: A.Experimental Response to RGS KO in yeast B.Response to RGS KO in the current model C.Reaction System with Selected Input(red) and Output(blue)

optimizing for reaction time (delay to steady-state) as well. It is possible to add s as a separate objective.

2 Delay vs. Efficiency Trade-off

Fig. 1C shows the example system, a submembrane compartment with a GPCR (G-protein coupled receptor) pathway from a mouse embryonic fibroblast, with ISO as input (extracellular ligand to beta-adrenergic receptors) and the phosphorylation of a membrane protein VASP as output. This model was implemented as an ODE model with 23 reactions and 10 total (initial) concentrations, and I/O transfer functions were derived as in [7]. The parameters were adapted to experimental biological data (not shown, [2]). In Fig. 2A the original ISO/pVASP transfer function is shown, which we use here for further optimization. It is clear that a complex signaling system may have several inputs, and a large number of outputs or target proteins. It is easy to extend the present discussion to optimize for multiple outputs in parallel, and a system could also be optimized for a number of I/O functions. In that case other measures, such as mutual information, may be employed as objectives. The basic principle will be explained here for a single I/O pathway. The system was trained for a signal distribution between 10nM and 1 μ M by adjusting only the kinetic rate parameters. Fig. 2A shows the I/O transfer function before and after adjustment, maximizing for $R_{C,S}$ and/or delay. We see that optimizing for the response coefficient alone shifts the function to the right to better cover the input range. Optimizing for the delay alone shifts it to the left, speeding up signals in the lower range (which are slower). This corresponds to the theoretical results in [7]. Both objectives together (equally weighted) are close to the original, biologically validated curve. The optimization method used is a simplex algorithm [6]. These results are simple, intuitive and encourage continuing to explore the basic idea.

3 Optimization and Evolutionary Learning

In principle, we may use any parameters in a system, concentrations or kinetic rates, to maximize signal transmission. For the evolutionary learning which produces a cellular system, it is the kinetic rates which are subject to evolution of protein structure and interactions, while concentrations may be regulated adaptively in each cell. Nonetheless there are expected concentration ranges for healthy cells, specific by cell type, which are the basis for kinetic rate optimization. During evolution, new protein subtypes develop with a different set of interactions. This corresponds to an adjustment of the available set of reactions, a type of structural learning to overcome the bottlenecks that are a result of tightly specified molecular kinetics.

In other words, we assume that kinetic rate learning operates on evolutionary time-scales, and that biologically attested signal transduction pathways contain sets of reactions and reaction rates which are optimal in terms of signal transmission efficiency. Our experiments show that kinetic rate solutions are not unique, yet there are clear tendencies on how to improve fitness of




Figure 2: A. Transfer Functions after Optimization of Kinetic Rates by $R_{pVASP,ISO}$, delay B. Relative Reaction Parameters during Optimization C. Binding Coefficients for High (red) and Low (blue) Fitness

a specific I/O signal transduction pathway (Fig. 2B). The initial activation of PKA by cAMP to form PKA2cAMP, the binding of ligand to receptor, the action of the GTPase RGS, the activity of PDE are all highly relevant, while phosphorylation of the receptor or of the cAMP synthesis enzyme AC are less relevant. This is evidenced by the degree of deviation of a reaction rate from the original as fitness improves (from left to right). (These are results from different experiments with different starting configurations, which were sorted according to R value. The delay values are not shown.) Again, this is encouraging, since it corresponds to accumulated biological observations on the relative importance of e.g. GTPases or PDE versus e.g. feedback phosphorylation by PKA.

Fig. 2C shows the distribution of binding parameters according to fitness value (R, delay) of the signal transduction pathway. We see a distinct clustering of binding coefficients, pointing to optimal ranges for these parameters. It is well-known that matching models to experimental time-series data yields sometimes large ranges of suitable parameters [8].

This tool can now be used in two different ways (a) in order to optimize signal transduction using experimental boundaries on the rate parameters as constraints (b) in order to measure the signal transmission efficiency in a biologically validated system [3] and compare it to what is theoretically possible given the set of equations ('motifs').

4 Co-regulation as an Adaptive Response

Most significantly, we now have a tool to investigate perturbations on concentrations. This is particularly interesting in the following sense: usually, when we use a computational model to investigate the effect of perturbation, we only study the effects. Now we can make the system itself adjust to the perturbation by utilizing optimization. We have created a self-regulating system, which uses objective functions to adapt to perturbation. This basic idea is extremely powerful, and could be used with different kinds of constraints on parameters, reaction times, etc. and with different I/O submodules of a larger system (such as cellular compartments). Here we are looking at a simple case (Fig. 3A): we reduce the level of a given protein (such as a kinase) to 10% of its usual concentration. As a result, all concentrations in the system are allowed to adjust until optimality of signal transmission and delay is reinstated. This process mimics co-regulation of protein expression in cellular systems, which is important in disease progression and often detrimental to targeted interventions. The model shows strong responses to reduction of the output protein VASP, and the phosphatase (PP1), which prevents pVASP (phosphorylated VASP) accumulation. Lowering PP1 lowers most other proteins, to re-establish sensitivity, lowering VASP increases most other concentrations to re-establish a clear signal. PDE4B and AC6 are co-regulated in the sense that both are reduced together. This is interesting because it concurs with the biological facts. AC6 synthesizes cAMP and PDE4B degrades it. I.e. the basic goal of the system is to provide for optimal signal transmission, here by homeostasis for the messenger molecule cAMP. Mostly it is only a small set of proteins that respond with altered concentrations to a single perturbation. This result confirms a view of signal transduction as highly modular [7]. Concentration changes may be




Figure 3: A. Response to Reduction to 10% of Original Concentration B. Transfer Function Response to Shifts in Input Signal Range C. Concentrations Changes in Response to Extracellular Signal Shift

caused by genetic up- or down- regulation, secretion and re-uptake, increased degradation, RNA interference, etc. and are therefore not easy to model from the standpoint of mechanistic biological modeling, which would need modules for all biologically attested processes. A unified perspective by a set of constraints and a set of objectives, such as has been envisaged here, may lead to better predictive results and may also be used as a guiding principle in constructing mechanistic models.

5 Optimal signal transmission depends on the signal

From the standpoint of disease modeling, an unusual protein concentration may be an adaptive response where a still functioning cell in a dysfunctional external signaling environment struggles to keep signal transmission efficient to support cellular function. In such a case, targeting this protein by pharmacological intervention will lead to co-regulation on other proteins. In general, as well as in real biological signaling systems, it is the localization of the input signaling range and the distribution of signals that the system transmits which are important for optimization. If we select the signal set that we optimize for in the right way, the input range will move towards the loglinear range, i.e. between the lower and upper input boundaries (Fig. 3B). As a result, a number of internal concentrations will change. Fig. 3C shows the relative concentration changes that result from a shift in input range. Interestingly, the low shift requires mostly a reduction in RGS. The high shift (which was also less effective) affects most concentrations, notably PKA. The results of this process may now be compared with biological evidence, e.g. for addiction, where this kind of "sensitization" is well-attested, but also for cancer where intercellular signaling is affected. Gene expression data may then be mined not only for evidence of the mechanics of genetic regulatory pathways but also for evidence of shifts in extracellular signaling, which cause altered protein expression.

6 Summary

The idea to look for parametric optimization as the basis of realistic cellular properties has been applied with success to metabolic fluxes [4]. In that case, optimization of growth is usually regarded as the single objective function. Here we use another objective function, signal transmission efficiency, to study the adaptive response of a signaling system to perturbation in concentrations. This equals maximization of concentration change in a target species in response to input signal. Optimization of growth and optimization of signal transmission are therefore related. We also optimize for the speed of the transmission which counteracts signal efficiency due to the basic properties of kinetic equations [7]. We use optimization in a two-step process: (a) in evolutionary learning, this is used to find kinetic rates for estimated concentrations (b) in cellular adaptation, this is used to re-calibrate the model in response to perturbations in concentrations or changes in extracellular signaling.

- [1] Bhalla, U and Iyengar, R. Emergent Properties of Networks of Biological Signaling Pathways. *Science*, 283(5400):381–387, 1999.
- [2] Blackman BE, Horner K, Heidmann J, Wang D, Richter W, Rich TC, and Conti M. PDE4D and PDE4B function in distinct subcellular compartments in mouse embryonic fibroblasts. *Journal of Biological Chemistry*, 286(14):12590–601, 2011.
- [3] Cheong R, Rhee A, Wang C, Nemenman I, and Levchenko A. Information transduction capacity of noisy biochemical signaling networks. *Science*, 334(6054):354–8, 2011.
- [4] Palsson BO, Edwards JS, Ibarra RU. In silico predictions of Escherichia coli metabolic capabilities are consistent with experimental data. *Nature Biotechnology*, 19(2):125–30, 2001.
- [5] Ferrell JE. Signaling motifs and Weber’s law. *Molecular Cell*, 36(5):724–7, 2009.
- [6] Teukolsky SA, Vetterling WT, Flannery BP, Press WH. Section 10.5. Downhill Simplex Method in Multidimensions. In: *Numerical Recipes: The Art of Sci. Computing*, 2007.
- [7] Scheler G. Transfer Functions for Protein Signal Transduction: Application to a Model of Striatal Neural Plasticity. *arxiv qbio 1208.1054*, 2012.
- [8] Secrier, M, Toni T and Stumpf M. The ABC of reverse engineering biological signalling systems. *Molecular BioSystems*, 5(12):1925–1935, 2009.
- [9] Yi TM, Kitano H, and Simon M. A quantitative characterization of the yeast heterotrimeric G protein cycle. *PNAS*, 100(19):10764–9, 2003.